

# Davood Fallahdar

MD&Surg (Hons), FRCS (Eng), Chir Plast (Hons), MicroSurg (Hons), ORL  
Consultant Plastic, Reconstructive & Aesthetic Surgeon



## POST-SURGERY INSTRUCTIONS: OTOPLASTY

Once your surgery is completed, you must follow all the instructions given to you in order to heal properly and have a good outcome.

The following instructions are your obligation. Use this as a checklist of progress as you heal. Included are normal post-surgical experiences and key health considerations that may be a cause of concern.

### TYPICAL POST-OPERATIVE SYMPTOMS

Typical symptoms and signs to watch for following otoplasty include:

**Swollen, red ears. A firm feeling or tingling of the outer ear.** These are normal experiences as the skin, tissues and sensory nerves heal. Pain medication will help you cope with any discomfort.

**Consistent sharp pain should be reported to our office immediately.**

**To alleviate any discomfort, and to reduce swelling, you may apply cool (not cold) compresses to your ears.** Do not apply ice or anything frozen directly on the skin. Soak soft plain white washcloths or gauze squares in ice water and wring out well. Apply any compress gently; do not apply any pressure, this could injure ears. Apply cool compresses for no longer than 20-minute intervals. Do not apply any heat.

### CALL THE OFFICE IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:

- **A high fever, severe nausea and vomiting, continued dizziness or incoherent behaviour, such as hallucinations.**
- **Any pain that cannot be controlled by your pain medication.**
- **Excessive bleeding or fluid seeping through the incisions.**
- **A severely misshapen appearance, excessive bruising or fluid retention that is localised to one side of the face or head.**

### DRESSING

You may be wearing a dressing, ace wrap or similar compression around your head to support your ears as they begin to heal in their new position and/or shape. Follow the directions you are given explicitly for wearing this dressing. **It is important that dressing not be removed or dislodged for any reason, except with explicit instructions from Mr. Fallahdar. Doing so may significantly impair the outcome of your procedure.**

### DAY OF SURGERY INSTRUCTIONS

You will only be released to the care of a responsible adult. All of these instructions must be clear to the adult who will monitor your health and support you.



BAPRAS

Mr Davood Fallahdar MD&Surg (Hons), FRCS (Eng), Chir Plast (Hons), MicroSurg (Hons), ORL GMC 4686602

Private Secretary: t 020 7352 8189 f 020 7352 8876 m 07506 660660 e info@davoodfallahdar.co.uk w www.davoodfallahdar.co.uk

BUPA Cromwell Hospital: 162-174 Cromwell Road, London SW5 0UT t 020 7460 5700

Cadogan Clinic: 120 Sloane Street, London SW1X 9BW t 020 7901 8500

# Davood Fallahdar

MD&Surg (Hons), FRCS (Eng), Chir Plast (Hons), MicroSurg (Hons), ORL  
Consultant Plastic, Reconstructive & Aesthetic Surgeon



\_\_\_\_\_ **Rest, but not bed rest.** While rest is important in the early stages of healing, equally important is that you are ambulatory: meaning that you are walking under your own strength. Spend 10 minutes every 2 hours engaged in light walking indoors as you recover.

\_\_\_\_\_ **Recline, do not lie down.** This will be more comfortable for you, and can reduce swelling. Always keep your head elevated. Do not bend forward or over.

\_\_\_\_\_ **Good nutrition.** Fluids are critical following surgery. Stick to cool, non-carbonated, non-alcoholic, caffeine-free and green tea-free beverages including fruit juices and water, milk and yogurt drinks. Stick with soft, bland, nutritious food for the first 24 hours.

\_\_\_\_\_ **Take all medication such as pain medication, antibiotics and other medication, exactly as prescribed.**

\_\_\_\_\_ **Wound care may be required.** You may be instructed to remove any dressing or compression to care for wounds. Do not remove any steri-strips or crusting near your stitches. Immediately replace any dressing or compression as directed.

\_\_\_\_\_ **Do not smoke.** Smoking can greatly impair your safety prior to surgery and your ability to heal following surgery. You must not smoke.

\_\_\_\_\_ **Relax.** Do not engage in any stressful activities. Take care of no one, including yourself. Let others tend to you.

## TWO TO SEVEN DAYS FOLLOWING SURGERY

During this time you will progress with each day that passes. Ease into your daily activities. You will receive clearance to begin return to school or work at your post-operative visit.

- **Practice daily sun protection.** An SPF 30 is essential regardless of the weather or your activities.
- **You may shower as needed. You must follow all the instructions you are given for compression, dressings and wound care.**
- **Apply skincare cautiously.** Do not use glycolics, retinoids or other potentially irritating skincare or haircare products until you receive clearance to do so.
- **Take antibiotic medications and supplements as directed.** Take pain medication only as needed. You may wish to switch from prescriptive pain medication to paracetamol or ibuprofen.
- **Continue to keep your head elevated, including when sleeping.**
- **Do not resume any exercise other than regular walking.** Walking is essential every day to prevent the formation of blood clots.
- **Maintain a healthy diet. Do not smoke. Do not consume alcohol.**



BAPRAS

Mr Davood Fallahdar MD&Surg (Hons), FRCS (Eng), Chir Plast (Hons), MicroSurg (Hons), ORL GMC 4686602

Private Secretary: t 020 7352 8189 f 020 7352 8876 m 07506 660660 e info@davoodfallahdar.co.uk w www.davoodfallahdar.co.uk

BUPA Cromwell Hospital: 162-174 Cromwell Road, London SW5 0UT t 020 7460 5700

Cadogan Clinic: 120 Sloane Street, London SW1X 9BW t 020 7901 8500

# Davood Fallahdar

MD&Surg (Hons), FRCS (Eng), Chir Plast (Hons), MicroSurg (Hons), ORL  
Consultant Plastic, Reconstructive & Aesthetic Surgeon

## ONE to FOUR WEEKS FOLLOWING SURGERY

As you resume your normal daily activities, you must continue proper care and healing.

- **Refrain from direct sun exposure.** If you are outdoors, apply at least an SPF 30 at least 30 minutes prior to sun exposure and wear a hat with a brim that shades your ears. Your skin is highly susceptible to sunburn or the formation of irregular, darkened pigmentation.
- **Do not smoke.** Smoking deprives your body of necessary oxygen that can result in poorly healed, wide, raised scars.
- **You may begin to sleep in a modified reclining position.** However do not sleep on your side, or with your head resting on your ear.
- **Do not engage in any contact sports or swimming until you are given clearance to do so.**

## SIX WEEKS FOLLOWING SURGERY

Healing will progress; swelling continues to diminish.

- **Discomfort or tightness and tingling of the ears will resolve.**
- **You may ease into your regular fitness routine.** However, if you are going to engage in any contact sports you must wear proper protection.
- **No need to resume smoking.** You have now gone 10 weeks (4 weeks prior to surgery and 6 weeks following) without a cigarette. For your long-term health, there is no need to resume smoking.

## YOUR FIRST YEAR

**Your appearance will change with age.** If your condition changes in any way, contact our office. Call us at any time with any of your questions or concerns.



Mr Davood Fallahdar MD&Surg (Hons), FRCS (Eng), Chir Plast (Hons), MicroSurg (Hons), ORL GMC 4686602

Private Secretary: t 020 7352 8189 f 020 7352 8876 m 07506 660660 e info@davoodfallahdar.co.uk w www.davoodfallahdar.co.uk

BUPA Cromwell Hospital: 162-174 Cromwell Road, London SW5 0UT t 020 7460 5700

Cadogan Clinic: 120 Sloane Street, London SW1X 9BW t 020 7901 8500